



## 2014 New CP<sup>3</sup>R Measures

### BREAST

- Radiation therapy is considered or administered within 1 year (365 days) of diagnosis for women undergoing mastectomy for breast cancer with  $\geq 4$  positive regional lymph nodes.
- Needle biopsy is performed prior to surgical treatment of breast cancer
- Breast conservation rate for women with AJCC stage 0, I, or II breast cancer (*surveillance measure*)

### NON-SMALL CELL LUNG

- Systemic chemotherapy is considered or administered within 4 months pre-operatively or 6 months post-operatively for surgically resected cases with pathologic lymph node positive (pN1/pN2) non-small cell lung cancer
- Surgery is not the *initial* treatment for patients with AJCC clinical N2 non-small cell lung cancer
- 10 or more lymph nodes are pathologically examined for resected non-small cell lung cancer (Stage I-II)
- Type of pulmonary resection for non-small cell lung cancer (*surveillance measure*)

### GI

- Chemotherapy is considered or administered within 4 months pre-operatively or 3 months post-operatively for AJCC stage IB – IV-M0 **gastric cancers**
- 15 or more lymph nodes are pathologically examined for resected **gastric cancers**
- Neo-adjuvant chemotherapy and radiation is considered or administered within 4 months pre-operatively for AJCC clinical stage T3-T4 N0 M0 or T1-T4 N1 M0 **esophageal cancer**
- **Esophageal** surgical volume (*surveillance measure*)
- **Pancreatic** surgical volume (*surveillance measure*)

#### **REVISION for Rectal cancer**

- Chemo/radiation is considered or administered within 3 months pre-operatively, or chemotherapy is considered or administered within 90 days post-operatively for AJCC stage II or III rectal cancer.